



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

JOURNAL OF
COMPUTATIONAL AND
APPLIED MATHEMATICS

Journal of Computational and Applied Mathematics 169 (2004) 345–357

www.elsevier.com/locate/cam

Two positive solutions for second-order quasilinear differential equation boundary value problems with sign changing nonlinearities[☆]

Yanping Guo^{a,b,*}, Jiwei Tian^a^a Marine Environment College, Ocean University of China, Qingdao 266003, PR China^b College of Sciences, Hebei University of Science and Technology, Shijiazhuang, Hebei 050018, PR China

Received 15 January 2003; received in revised form 1 October 2003

Abstract

In this paper, the second order quasilinear differential equation $(\Phi(y'))' + q(t)f(t, y) = 0$, $0 < t < 1$ subject to Dirichlet boundary conditions and mixed boundary conditions is studied, where f is allowed to change sign, $\Phi(v) = |v|^{p-2}v$, $p > 1$. We show the existence of at least two positive solutions by using a new fixed point theorem in cones.

© 2004 Elsevier B.V. All rights reserved.

Keywords: Quasilinear differential equation; Positive symmetric solution; Fixed point theorem in cones

1. Introduction

In this paper we shall consider the Dirichlet boundary value problem with a p -Laplacian operator

$$\begin{aligned}(\Phi(y'))' + q(t)f(t, y) &= 0, \quad 0 < t < 1, \\ y(0) = y(1) &= 0\end{aligned}\tag{1}$$

[☆] The project is supported by the Natural Science Foundation of China (10371030) and the Natural Science Foundation of Hebei Province (603384) and the Scientific Research Foundation of the Education Department of Hebei Province (2003216).

* Corresponding author.

E-mail address: guoyanping65@sohu.com (Y. Guo).

and the mixed boundary value problem with a p -Laplacian operator

$$\begin{aligned}(\Phi(y'))' + q(t)f(t, y) &= 0, \quad 0 < t < 1, \\ y(0) = y'(1) &= 0,\end{aligned}\tag{2}$$

where our nonlinear term f is allowed to change sign and $\Phi(v) = |v|^{p-2}v$, $p > 1$. The following conditions will be assumed throughout:

- (H₁) $f : [0, 1] \times [0, \infty) \rightarrow [-M, \infty)$ is continuous and $f(t, 0) \geq 0 (\neq 0)$ for $t \in [0, 1]$, where $M > 0$ is a constant;
 (H₂) $q : (0, 1) \rightarrow [0, \infty)$ is continuous and $q(t) \neq 0$ on any subinterval of $(0, 1)$, and $\int_0^1 q(t) dt < \infty$.

In the recent years, there has been increasing interest in questions of positive solutions of boundary value problems for ordinary differential equations [1,2,4,5,8,9,12,14], as well as for finite difference equations. The recent book by Agarwal et al. [1] gives a good overview of much of the work which has been done and the methods used.

In [9], Erbe and Wang studied the existence of positive solutions of the equation $u'' + q(t)f(u) = 0$ by using the Krasnosel'skii fixed point theorem [10], where $q(t)$ is continuous on $[0, 1]$ and $f(u)$ is continuous on $[0, \infty)$. Since then, the Krasnosel'skii fixed point theorem has been widely used to discuss the existence of positive solutions for boundary value problems. For the second-order boundary value problem

$$\begin{aligned}y'' + f(y) &= 0, \quad 0 \leq t \leq 1, \\ y(0) = y(1) &= 0,\end{aligned}\tag{3}$$

where $f : R \rightarrow [0, \infty)$ is continuous, Avery [2] imposed conditions on f to yield at least three symmetric positive solutions to (3) by applying the Leggett–Williams fixed point theorem [13]. Avery and Henderson [5] improved Avery's results by using a five functionals fixed point theorem [3]. On the other hand, Avery et al. [4] applied a twin fixed point theorem to obtain at least two positive solutions for the right focal boundary value problem

$$\begin{aligned}y'' + f(y) &= 0, \quad 0 \leq t \leq 1, \\ y(0) = y'(1) &= 0,\end{aligned}\tag{4}$$

where $f : R \rightarrow [0, \infty)$ is continuous.

Using the fixed point theorem in cones due to Krasnosel'skii, Junyu Wang [14] studied the existence of positive solutions of the quasilinear differential equation

$$(g(u'))' + q(t)f(u) = 0, \quad 0 < t < 1$$

subject to nonlinear boundary conditions, where $g(v) = |v|^{p-2}v$, $p > 1$.

In order to apply the concavity of solutions in the proofs, all the above results were done under the assumption that function f is nonnegative. For the sign changing nonlinearity f , few results were done. Motivated by the results mentioned above, in this paper we apply a new fixed point theorem in cones to obtain at least two positive solutions of problems (1) and (2). Growth conditions are imposed on f in later sections so that our fixed point theorem in cones is applicable.

2. The fixed point theorem in cones

For a cone K in a Banach space X with norm $\|\cdot\|$ and a constant $r > 0$, let $K_r = \{x \in K : \|x\| < r\}$, $\partial K_r = \{x \in K : \|x\| = r\}$. Suppose $\alpha : K \rightarrow R^+$ is a continuous functional, let

$$K(b) = \{x \in K : \alpha(x) < b\}, \partial K(b) = \{x \in K : \alpha(x) = b\}$$

and $K_a(b) = \{x \in K : a < \|x\|, \alpha(x) < b\}$. The origin in X is denoted by θ .

Definition 1. Given a cone K in a real Banach space X , a functional $\alpha : K \rightarrow R$ is said to be concave functional on K provided

$$\alpha(tx + (1-t)y) \geq t\alpha(x) + (1-t)\alpha(y)$$

for all $x, y \in K$ and $0 \leq t \leq 1$.

Theorem 1. Let X be a real Banach space with norm $\|\cdot\|$ and $K \subset X$ a cone. Suppose $T : K \rightarrow K$ is a completely continuous operator, and $\alpha : K \rightarrow R^+$ a continuous concave functional satisfying $\alpha(x) \leq \|x\|$ for all $x \in K$. If there are constants $c > b > a > 0$ such that

- (C₁) $\|Tx\| < a$ for $x \in \partial K_a$;
- (C₂) $\alpha(Tx) > b$ for $x \in \partial K(b)$;
- (C₃) $\alpha(x) \leq b$ implies $\|Tx\| < c$,

then T has in K at least a fixed points y such that

$$a < \|y\| < c, \quad \alpha(y) < b.$$

Proof. Let the symbol \deg_K denote the degree on cone K [6]. Then condition (C₁) implies

$$\deg_K \{I - T, K_a, \theta\} = 1.$$

Let $\Omega = K(b) \cap K_{2c}$, we now prove that

$$\deg_K \{I - T, \Omega, \theta\} = 0.$$

Conditions (C₂) and (C₃) imply $\inf_{x \in \partial K(b)} \alpha(Tx) \geq b > 0$ and $\inf_{x \in \partial K(b)} \|Tx\| \leq c$. Let $\tilde{T} : \bar{K}(b) \rightarrow K$ be an extension of $T|_{\partial K(b)} : \partial K(b) \rightarrow K$. Dugundji extension theorem [7] ensures that \tilde{T} is completely continuous and $\tilde{T}(\bar{K}(b)) \subset \overline{\text{conv}} T(\partial K(b))$. Since $\{x \in K : \alpha(x) \geq b\} \cap \{x \in K : \|x\| \leq c\}$ is a convex set, we have

$$\inf_{x \in \bar{K}(b)} \alpha(\tilde{T}x) \geq b > 0 \quad \text{and} \quad \inf_{x \in \bar{K}(b)} \|\tilde{T}x\| \leq c.$$

We claim

$$\deg_K \{I - \tilde{T}, \Omega, \theta\} = 0.$$

Clearly $\partial\Omega = (\partial K(b) \cap \bar{K}_{2c}) \cup (\partial K_{2c} \cap \bar{K}(b))$. For $x \in \partial\Omega$, $(I - \tilde{T})(x) \neq \theta$. If it is not true, then there exists $x_0 \in \partial\Omega$ such that

$$x_0 = \tilde{T}x_0.$$

If $x_0 \in \partial K(b) \cap \bar{K}_{2c}$, then

$$b = \alpha(x_0) = \alpha(\tilde{T}x_0) = \alpha(Tx_0) > b,$$

a contradiction. On the other hand, if $x_0 \in \partial K_{2c} \cap \bar{K}(b)$, then

$$2c = \|x_0\| = \|\tilde{T}x_0\| \leq c,$$

a contradiction. For $x \in \Omega$, we have

$$\alpha(x) < b \quad \text{and} \quad \alpha(\tilde{T}x) \geq b.$$

Thus, $(I - \tilde{T})(x) \neq \theta$ for $x \in \Omega$. It follows that

$$\deg_K\{I - \tilde{T}, \Omega, \theta\} = 0.$$

Take a homotopy $H(x, \lambda) = \lambda Tx + (1 - \lambda)\tilde{T}x$. It is easy to see that

$$H(x, \lambda) \neq x \quad \text{for all } x \in \partial\Omega, \lambda \in [0, 1].$$

Thus,

$$\deg_K\{I - T, \Omega, \theta\} = \deg_K\{I - \tilde{T}, \Omega, \theta\} = 0.$$

From $\alpha(x) \leq \|x\|$, we have $K_a \subset K(b) \cap K_{2c} = \Omega$. Then

$$\begin{aligned} \deg_K\{I - T, \Omega \setminus K_a, \theta\} \\ = \deg_K\{I - T, \Omega, \theta\} - \deg_K\{I - T, K_a, \theta\} \\ = -1. \end{aligned}$$

So T has in K a fixed point y such that

$$a < \|y\| < c, \quad \alpha(y) < b.$$

Theorem 1 is now proved. \square

3. Dirichlet boundary problem

In this section, we impose growth conditions on f which allow us to apply Theorem 1 to obtain two positive solutions of (1). Let $X = C[0, 1]$, $K = \{x \in X : x(t) \geq 0, x(t) = x(1 - t) \text{ for } t \in [0, 1]\}$, $K' = \{x \in K : x \text{ is concave on } [\delta/2, 1 - \delta/2]\}$, where $\delta \in (0, \frac{1}{2})$. Obviously, $K, K' \subset X$ are two cones with $K' \subset K$.

We suppose the following conditions are satisfied:

(H₃) $f(t, x), q(t)$ are symmetric about $t = \frac{1}{2}$;

and there are $a, b, d > 0$ satisfying

$$0 < \frac{1 - \delta}{2} \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) dt \right] + d \leq a < \delta b < b$$

such that

$$(H_4) \quad f(t, x) \int_{\delta/2}^{1/2} q(t) dt \geq M \int_0^{\delta/2} q(t) dt \quad \text{for } (t, x) \in [\delta/2, 1 - \delta/2] \times [d, b];$$

$$(H_5) \quad f(t, x) \int_0^{1/2} q(t) dt < \Phi(2a) \quad \text{for } (t, x) \in [0, 1] \times [0, a];$$

$$(H_6) \quad f(t, x) \int_{\delta}^{1/2} q(t) dt \geq M \int_0^{\delta} q(t) dt + \Phi(b) \quad \text{for } (t, x) \in [\delta, 1 - \delta] \times [\delta b, b];$$

$$(H_7) \quad f(t, x) \int_0^{\delta/2} q(t) dt \leq \Phi(2a/\delta) \quad \text{for } (t, x) \in [0, \delta/2] \times [0, b].$$

For $x \in K$ we define

$$\alpha(x) = \min_{\delta \leq t \leq 1-\delta} x(t)$$

$$(Tx)(t) = \begin{cases} \left(\int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) F(u, x(u)) du \right) ds \right)^+, & 0 \leq t \leq \frac{1}{2}, \\ \left(\int_t^1 \Phi^{-1} \left(\int_{1/2}^s q(u) F(u, x(u)) du \right) ds \right)^+, & \frac{1}{2} \leq t \leq 1, \end{cases}$$

where $(B)^+ = \max\{B, 0\}$, $F(t, u) = f(t, u)$ for $u \geq 0$ and $F(t, u) = f(t, 0)$ for $u < 0$.

$$(Ax)(t) = \begin{cases} \int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) F(u, x(u)) du \right) ds, & 0 \leq t \leq \frac{1}{2}, \\ \int_t^1 \Phi^{-1} \left(\int_{1/2}^s q(u) F(u, x(u)) du \right) ds, & \frac{1}{2} \leq t \leq 1. \end{cases}$$

For $x \in X$, define $\theta : X \rightarrow K$ by $(\theta x)(t) = \max\{x(t), 0\}$, then $T = \theta \circ A$. For $x \in K'$, let

$$(T'x)(t) = \begin{cases} \int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) f^*(u, x(u)) du \right) ds, & 0 \leq t \leq \frac{1}{2}, \\ \int_t^1 \Phi^{-1} \left(\int_{1/2}^s q(u) f^*(u, x(u)) du \right) ds, & \frac{1}{2} \leq t \leq 1, \end{cases}$$

where

$$f^*(t, x) = \begin{cases} f(t, 0), & 0 \leq t < \frac{\delta}{2} \text{ or } 1 - \frac{\delta}{2} < t \leq 1, -\infty < x < 0, \\ f(t, x), & \frac{\delta}{2} \leq t \leq 1 - \frac{\delta}{2}, d \leq x \leq b, \\ f(t, d), & \frac{\delta}{2} \leq t \leq 1 - \frac{\delta}{2}, -\infty < x < d, \\ f(t, x), & 0 \leq t < \frac{\delta}{2} \text{ or } 1 - \frac{\delta}{2} < t \leq 1, 0 \leq x \leq b, \\ f(t, b), & 0 \leq t \leq 1, b < x < +\infty. \end{cases}$$

Lemma 1 (Guo Yanping et al. [11]). Suppose $A : K \rightarrow X$ is completely continuous, then $\theta \circ A : K \rightarrow K$ is also completely continuous.

Condition (H_3) implies that A and T' are well defined. From the continuity of f , it is easy to see that $A : K \rightarrow X$ is completely continuous. So $T : K \rightarrow K$ is completely continuous by using Lemma 1. For $x \in K'$, we have $x(t) \geq \delta/(1-\delta) \max_{\delta/2 \leq t \leq 1-\delta/2} x(t) \geq \delta \max_{\delta/2 \leq t \leq 1-\delta/2} x(t)$ for $t \in [\delta, 1-\delta]$ by the concavity of x on $[\delta/2, 1-\delta/2]$. Thus, $\alpha(x) \leq \|x\|$ and $\alpha(x) \geq \delta \max_{\delta/2 \leq t \leq 1-\delta/2} x(t)$.

Lemma 2. Suppose (H_1) – (H_4) hold. Then $T' : K' \rightarrow K'$ is completely continuous.

Proof. For all $x \in K'$, from (H_1) and (H_4) , we have

$$\begin{aligned} \int_s^{1/2} q(u) f^*(u, x(u)) du &= \int_s^{\delta/2} q(u) f^*(u, x(u)) du + \int_{\delta/2}^{1/2} q(u) f^*(u, x(u)) du \\ &\geq -M \int_0^{\delta/2} q(u) du + \int_{\delta/2}^{1/2} q(u) \cdot \frac{M \int_0^{\delta/2} q(t) dt}{\int_{\delta/2}^{1/2} q(t) dt} du \\ &= 0 \text{ for } 0 \leq s \leq \frac{\delta}{2}, \\ \int_s^{1/2} q(u) f^*(u, x(u)) du &\geq 0 \text{ for } \frac{\delta}{2} \leq s \leq \frac{1}{2}, \end{aligned}$$

thus,

$$\begin{aligned} (T'x)(t) &= \int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) f^*(u, x(u)) du \right) ds \geq 0 \text{ for } 0 \leq t \leq \frac{1}{2}, \\ (\Phi((T'x)'))'(t) &= -q(t) f^*(t, x(t)) \leq 0 \text{ for } t \in \left[\frac{\delta}{2}, 1 - \frac{\delta}{2} \right]. \end{aligned}$$

So $T' : K' \rightarrow K'$. Using the continuity of f and the definition of f^* , it is easy to see that $T' : K \rightarrow K'$ is completely continuous. \square

Theorem 2. Suppose (H_1) – (H_7) hold. Then the boundary value problem (1) has at least two positive solutions y_1 and y_2 such that

$$0 < \|y_1\| < a < \|y_2\|, \min_{t \in [\delta, 1-\delta]} y_2(t) < \delta b.$$

Proof. At first we show that T has a fixed point $y_1 \in K$ with $0 < \|y_1\| < a$. In fact, for all $x \in \partial K_a$, we have $\|x\| = a$. From (H_5) we obtain

$$\begin{aligned} \|Tx\| &= \max_{0 \leq t \leq 1/2} \left| \left[\int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) f(u, x(u)) du \right) ds \right]^+ \right| \\ &= \max_{0 \leq t \leq 1/2} \max \left\{ \int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) f(u, x(u)) du \right) ds, 0 \right\} \end{aligned}$$

$$\begin{aligned}
&< \frac{1}{2} \Phi^{-1} \left(\int_0^{1/2} q(u) \cdot \frac{1}{\int_0^{1/2} q(t) dt} \Phi(2a) du \right) \\
&= \frac{1}{2} \cdot 2a = a.
\end{aligned}$$

The existence of y_1 is proved by using the Schauder fixed point theorem.

Obviously, y_1 is a solution of (1) if and only if y_1 is a fixed point of A . Next we will prove that y_1 is a fixed point of A . Suppose this is not true, then there is $t_0 \in (0, 1)$ such that $y_1(t_0) \neq (Ay_1)(t_0)$. It must be $(Ay_1)(t_0) < 0 = y_1(t_0)$. Let (t_1, t_2) be the interval such that $t_0 \in (t_1, t_2)$, $(Ay_1)(t) < 0$ for all $t \in (t_1, t_2)$, and $(Ay_1)(t_1) = (Ay_1)(t_2) = 0$. It is easy to see that $[t_1, t_2] \neq [0, 1]$ by (H_1) . Without loss of generality, suppose $t_2 < 1$. Then $y_1(t) \equiv 0$ for $t \in [t_1, t_2]$, and $(Ay_1)(t) < 0$ for $t \in (t_1, t_2)$, and $(Ay_1)(t_2) = 0$. Thus, $(Ay_1)'(t_2) \geq 0$. (H_1) implies $(\Phi((Ay_1)'))'(t) = -q(t)f(t, 0) \leq 0$ for $t \in (t_1, t_2)$. So, $(Ay_1)'(t) \geq 0$ for $t \in [t_1, t_2]$. Therefore, $t_1 = 0$. We get $(Ay_1)(0) \leq (Ay_1)(t_0) < 0$, a contradiction.

We now show that (C_1) of Theorem 1 is satisfied. For $x \in \partial K'_a$, we have $\|x\| = a$. From (H_5) we obtain

$$\begin{aligned}
\|T'x\| &= \max_{0 \leq t \leq 1/2} \left| \int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) f^*(u, x(u)) du \right) ds \right| \\
&< \frac{1}{2} \Phi^{-1} \left(\int_0^{1/2} q(u) \cdot \frac{1}{\int_0^{1/2} q(t) dt} \Phi(2a) du \right) \\
&= \frac{1}{2} \cdot 2a = a.
\end{aligned}$$

Next, we show that (C_2) of Theorem 1 is satisfied. For $x \in \partial K'(\delta b)$, i.e., $\alpha(x) = \delta b$. For $\delta \leq t \leq 1 - \delta$, we have $\delta b \leq x(t) \leq b$. we use conditions (H_1) and (H_6) to obtain

$$\begin{aligned}
\alpha(T'x) &= \min_{\delta \leq t \leq 1/2} \int_0^t \Phi^{-1} \left(\int_s^{1/2} q(u) f^*(u, x(u)) du \right) ds \\
&= \int_0^\delta \Phi^{-1} \left(\int_s^{1/2} q(u) f^*(u, x(u)) du \right) ds \\
&> \delta \Phi^{-1} \left(-M \int_0^\delta q(u) du + \int_\delta^{1/2} q(u) \cdot \frac{M \int_0^\delta q(t) dt + \Phi(b)}{\int_\delta^{1/2} q(t) dt} du \right) \\
&= \delta b.
\end{aligned}$$

Using the continuity of f and the definition of f^* , there is $c > b$ such that $\|T'x\| < c$ for $\alpha(x) \leq b$. Applying Theorem 1, T' has a fixed point y_2 such that $y_2 \in K'_a(\delta b)$.

Finally, we show that $Ax = T'x$ for $x \in K'_a(\delta b) \cap \{u : T'u = u\}$. Let $x \in K'_a(\delta b) \cap \{u : T'u = u\}$, then

$$\|x\| > a \geq \frac{1-\delta}{2} \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) dt \right] + d.$$

We claim $\|x\| = \max_{\delta/2 \leq t \leq 1-\delta/2} x(t)$. If there is $t_0 \in (0, \delta/2)$ such that $x(t_0) = \|x\| > a$, then

$$x'(t_0) = (A'x)'(t_0) = \Phi^{-1} \left(\int_{t_0}^{1/2} q(u) f^*(u, x(u)) du \right) = 0,$$

i.e.,

$$\int_{t_0}^{1/2} q(u) f^*(u, x(u)) du = 0.$$

From (H₇), we have

$$\begin{aligned} x(t_0) = \|x\| &= \int_0^{t_0} \Phi^{-1} \left(\int_s^{1/2} q(u) f^*(u, x(u)) du \right) ds \\ &= \int_0^{t_0} \Phi^{-1} \left(\int_s^{t_0} q(u) f^*(u, x(u)) du \right) ds \\ &\leq \frac{\delta}{2} \Phi^{-1} \left(\int_0^{\delta/2} q(u) \cdot \frac{1}{\int_0^{\delta/2} q(t) dt} \Phi \left(\frac{2a}{\delta} \right) du \right) \\ &= a, \end{aligned}$$

a contradiction. Therefore, $\|x\| = \max_{\delta/2 \leq t \leq 1-\delta/2} x(t)$.

Next we will prove $x(\delta/2) \geq d$. Suppose this is not true, then there exists $t_0 \in (\delta/2, \frac{1}{2})$ such that

$$x'(t_0) > \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) dt \right].$$

It follows from the concavity of x on $[\delta/2, 1 - \delta/2]$ that

$$x \left(\frac{\delta}{2} \right) \geq x'(t_0) > \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) dt \right].$$

For $0 \leq t \leq \delta/2$, we have

$$\begin{aligned} \Phi(x'(t)) &= \Phi \left[x' \left(\frac{\delta}{2} \right) \right] - \int_t^{\delta/2} (\Phi(x'(s)))' ds \\ &= \Phi \left[x' \left(\frac{\delta}{2} \right) \right] + \int_t^{\delta/2} q(s) f^*(s, x(s)) ds \\ &\geq \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) dt \right] - M \int_0^{\delta/2} q(s) ds \\ &= \Phi \left(\frac{2d}{\delta} \right), \end{aligned}$$

i.e., $x'(t) \geq 2d/\delta$. Therefore,

$$0 = x(0) = x \left(\frac{\delta}{2} \right) - \int_0^{\delta/2} x'(s) ds < d - \frac{\delta}{2} \cdot \frac{2d}{\delta} = 0,$$

a contradiction. Thus, $d \leq x(t) \leq b$ for $\delta/2 \leq t \leq 1 - \delta/2$. From the definition of f^* , we have $f^*(t, x(t)) = f(t, x(t))$ for $0 \leq t \leq 1$. Then $Ax = T'x$ for $x \in K'_a(\delta b) \cap \{u : T'u = u\}$. Thus, y_2 is a solution of (1). Theorem 2 is now completed. \square

4. Mixed boundary problem

Let $X = C[0, 1]$, $P = \{x \in X : x(t) \geq 0 \text{ for } t \in [0, 1]\}$, $P' = \{x \in P : x \text{ is concave, nondecreasing on } [\delta/2, 1]\}$, where $\delta \in (0, 1)$. Obviously, $P, P' \subset X$ are two cones with $P' \subset P$.

We suppose that there are $a, b, d > 0$ satisfying

$$0 < \frac{2-\delta}{2} \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) dt \right] + d \leq a < \frac{\delta}{2-\delta} b < b$$

such that

$$(G_1) \quad f(t, x) \int_{\delta/2}^1 q(t) dt \geq M \int_0^{\delta/2} q(t) dt \text{ for } (t, x) \in [\delta/2, 1] \times [d, b];$$

$$(G_2) \quad f(t, x) \int_0^1 q(t) dt < \Phi(a) \text{ for } (t, x) \in [0, 1] \times [0, a];$$

$$(G_3) \quad f(t, x) \int_{\delta}^1 q(t) dt \geq M \int_0^{\delta} q(t) dt + \Phi(b/(2-\delta)) \text{ for } (t, x) \in [\delta, 1] \times [\delta/(2-\delta)b, b];$$

$$(G_4) \quad f(t, x) \int_0^{\delta/2} q(t) dt \leq \Phi \left(\frac{2a}{\delta} \right) \text{ for } (t, x) \in [0, \delta/2] \times [0, b].$$

For $x \in P$ we define

$$\alpha(x) = \min_{\delta \leq t \leq 1} x(t);$$

$$(T_1 x)(t) = \left[\int_0^t \Phi^{-1} \left(\int_s^1 q(u) F(u, x(u)) du \right) ds \right]^+;$$

$$(A_1 x)(t) = \int_0^t \Phi^{-1} \left(\int_s^1 q(u) F(u, x(u)) du \right) ds;$$

$$(T'_1 x)(t) = \int_0^t \Phi^{-1} \left(\int_s^1 q(u) f^*(u, x(u)) du \right) ds,$$

where

$$f^*(t, x) = \begin{cases} f(t, 0), & 0 \leq t < \frac{\delta}{2}, -\infty < x < 0, \\ f(t, x), & \frac{\delta}{2} \leq t \leq 1, d \leq x \leq b, \\ f(t, d), & \frac{\delta}{2} \leq t \leq 1, -\infty < x < d, \\ f(t, x), & 0 \leq t < \frac{\delta}{2}, 0 \leq x \leq b, \\ f(t, b), & 0 \leq t \leq 1, b < x < +\infty. \end{cases}$$

Essentially the same reasoning as in Lemma 2, we can obtain $T_1 : P \rightarrow P$ and $T'_1 : P' \rightarrow P'$ are completely continuous. For $x \in P'$, we have $x(t) \geq \delta/(2-\delta) \max_{\delta/2 \leq t \leq 1} x(t)$ for $t \in [\delta, 1]$ by the concavity of x on $[\delta/2, 1]$. Thus, $\alpha(x) \leq \|x\|$ and $\alpha(x) \geq \delta/(2-\delta) \max_{\delta/2 \leq t \leq 1} x(t)$.

Theorem 3. Suppose (H_1) – (H_2) , (G_1) – (G_4) hold. Then the boundary value problem (2) has at least two positive solutions y_1 and y_2 such that:

$$0 < \|y_1\| < a < \|y_2\|, \min_{t \in [\delta, 1]} y_2(t) < \frac{\delta}{2-\delta} b.$$

Proof. At first we show that T_1 has a fixed point $y_1 \in P$ with $0 < \|y_1\| < a$. In fact, for all $x \in \partial P_a$, we have $\|x\| = a$. From (G_2) we obtain

$$\begin{aligned} \|T_1 x\| &= \max_{0 \leq t \leq 1} \left| \left[\int_0^t \Phi^{-1} \left(\int_s^1 q(u) F(u, x(u)) du \right) ds \right]^+ \right| \\ &< \Phi^{-1} \left(\int_0^1 q(u) \cdot \frac{1}{\int_0^1 q(t) dt} \Phi(a) du \right) = a. \end{aligned}$$

The existence of y_1 is proved by using the Schauder fixed point theorem.

Next we will prove that y_1 is a fixed point of A_1 . Suppose the contrary, then there exists $t_0 \in (0, 1)$ such that $y_1(t_0) \neq (A_1 y_1)(t_0)$. It must be $(A_1 y_1)(t_0) < 0 = y_1(t_0)$. Let (t_1, t_2) be the maximal interval such that $t_0 \in (t_1, t_2)$, $(A_1 y_1)(t) < 0$ for all $t \in (t_1, t_2)$. It is easy to see that $[t_1, t_2] \neq [0, 1]$ by (H_1) . If $t_2 < 1$, then $y_1(t) \equiv 0$ for $t \in [t_1, t_2]$, and $(A_1 y_1)(t) < 0$ for $t \in (t_1, t_2)$, and $(A_1 y_1)(t_2) = 0$. Thus, $(A_1 y_1)'(t_2) \geq 0$. (H_1) implies $(\Phi((A_1 y_1)'))'(t) = -q(t)f(t, 0) \leq 0$ for $t \in (t_1, t_2)$. So, $(A_1 y_1)'(t) \geq 0$ for $t \in [t_1, t_2]$. We obtain $t_1 = 0$. Therefore, $(A_1 y_1)(0) \leq (A_1 y_1)(t_0) < 0$, a contradiction. If $t_1 > 0$, then $y_1(t) \equiv 0$ for $t \in [t_1, t_2]$, and $(A_1 y_1)(t) < 0$ for $t \in (t_1, t_2)$, and $(A_1 y_1)(t_1) = 0$. Thus, $(A_1 y_1)'(t_1) \leq 0$. (H_1) implies $(\Phi((A_1 y_1)'))'(t) = -q(t)f(t, 0) \leq 0$ for $t \in (t_1, t_2)$. So, $(A_1 y_1)'(t) \leq 0$ for $t \in [t_1, t_2]$. We obtain $t_2 = 1$. On the other hand, $(A_1 y_1)'(1) = 0$, a contradiction.

We now show that (C_1) of Theorem 1 is satisfied. For $x \in \partial P'_a$, we have $\|x\| = a$. From (G_2) we obtain

$$\begin{aligned} \|T'_1 x\| &= \max_{0 \leq t \leq 1} \left| \int_0^t \Phi^{-1} \left(\int_s^1 q(u) f^*(u, x(u)) du \right) ds \right| \\ &< \Phi^{-1} \left(\int_0^1 q(u) \cdot \frac{1}{\int_0^1 q(t) dt} \Phi(a) du \right) = a. \end{aligned}$$

Next we show that (C_2) of Theorem 1 is satisfied. For $x \in \partial P'(\delta/(2-\delta)b)$, i.e., $\alpha(x) = \delta/(2-\delta)b$. For $\delta \leq t \leq 1$, we have $\delta/(2-\delta)b \leq x(t) \leq b$. We use conditions (H_1) and (G_3) to obtain

$$\begin{aligned} \alpha(T'_1 x) &= \min_{\delta \leq t \leq 1} \int_0^t \Phi^{-1} \left(\int_s^1 q(u) f^*(u, x(u)) du \right) ds \\ &= \int_0^\delta \Phi^{-1} \left(\int_s^1 q(u) f^*(u, x(u)) du \right) ds \end{aligned}$$

$$\begin{aligned}
&> \delta \Phi^{-1} \left[-M \int_0^\delta q(u) \, du + \int_\delta^1 q(u) \cdot \frac{M \int_0^\delta q(t) \, dt + \Phi(b/(2-\delta))}{\int_\delta^1 q(t) \, dt} \, du \right] \\
&= \frac{\delta}{2-\delta} b.
\end{aligned}$$

Using the continuity of f and the definition of f^* , there is $c > b$ such that $\|T'_1 x\| < c$ for $\alpha(x) \leq b$. Applying Theorem 1, T'_1 has a fixed point y_2 such that $y_2 \in P'_a(\delta/(2-\delta)b)$.

Finally, we show that $A_1 x = T'_1 x$ for $x \in P'_a(\delta/(2-\delta)b) \cap \{u : T'u = u\}$. Let $x \in P'_a(\delta/(2-\delta)b) \cap \{u : T'u = u\}$, then

$$\|x\| > a \geq \frac{2-\delta}{2} \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) \, dt \right] + d.$$

We claim $\|x\| = \max_{\delta/2 \leq t \leq 1} x(t)$. If there is $t_0 \in (0, \delta/2)$ such that $x(t_0) = \|x\| > a$, then

$$x'(t_0) = (A'_1 x)'(t_0) = \Phi^{-1} \left(\int_{t_0}^1 q(u) f^*(u, x(u)) \, du \right) = 0,$$

i.e.,

$$\int_{t_0}^1 q(u) f^*(u, x(u)) \, du = 0.$$

From (G₄), we have

$$\begin{aligned}
x(t_0) = \|x\| &= \int_0^{t_0} \Phi^{-1} \left(\int_s^1 q(u) f^*(u, x(u)) \, du \right) \, ds \\
&= \int_0^{t_0} \Phi^{-1} \left(\int_s^{t_0} q(u) f^*(u, x(u)) \, du \right) \, ds \\
&\leq \frac{\delta}{2} \Phi^{-1} \left[\int_0^{\delta/2} q(u) \cdot \frac{1}{\int_0^{\delta/2} q(t) \, dt} \Phi \left(\frac{2a}{\delta} \right) \, du \right] \\
&= a,
\end{aligned}$$

a contradiction. Therefore, $\|x\| = \max_{\delta/2 \leq t \leq 1} x(t)$.

Next we will prove $x(\delta/2) \geq d$. Suppose this is not true, then there exists $t_0 \in (\delta/2, 1)$ such that

$$x'(t_0) > \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) \, dt \right].$$

It follows from the concavity of x on $[\delta/2, 1]$ that

$$x' \left(\frac{\delta}{2} \right) \geq x'(t_0) > \Phi^{-1} \left[\Phi \left(\frac{2d}{\delta} \right) + M \int_0^{\delta/2} q(t) \, dt \right].$$

For $0 \leq t \leq \delta/2$, we have

$$\begin{aligned}\Phi(x'(t)) &= \Phi\left[x'\left(\frac{\delta}{2}\right)\right] - \int_t^{\delta/2} (\Phi(x'(s)))' ds \\ &= \Phi\left[x'\left(\frac{\delta}{2}\right)\right] + \int_t^{\delta/2} q(s)f^*(s, x(s)) ds \\ &\geq \left[\Phi\left(\frac{2d}{\delta}\right) + M \int_0^{\delta/2} q(t) dt\right] - M \int_0^{\delta/2} q(s) ds \\ &= \Phi\left(\frac{2d}{\delta}\right),\end{aligned}$$

i.e., $x'(t) \geq 2d/\delta$. Therefore,

$$0 = x(0) = x\left(\frac{\delta}{2}\right) - \int_0^{\delta/2} x'(s) ds < d - \frac{\delta}{2} \cdot \frac{2d}{\delta} = 0,$$

a contradiction. Thus, $d \leq x(t) \leq b$ for $\delta/2 \leq t \leq 1$. From the definition of f^* , we have $f^*(t, x(t)) = f(t, x(t))$ for $0 \leq t \leq 1$. Then $A_1x = T'_1x$ for $x \in P'_a(\delta/(2-\delta)b) \cap \{u : T'_1u = u\}$. So, y_2 is a solution of (2). Thus, the problem (2) has at least two positive solutions y_1 and y_2 such that

$$0 < \|y_1\| < a < \|y_2\|, \min_{t \in [\delta, 1]} y_2(t) < \frac{\delta}{2-\delta} b. \quad \square$$

Example. In problem (1), suppose that $p = 3$, $q(t) = 1$, and

$$f(t, x) = \begin{cases} 1 - 16x^2, & 0 \leq t \leq 1, 0 \leq x < 1/2, \\ -7 + 8x, & 0 \leq t \leq 1, 1/2 \leq x < 1, \\ 1 + 10(x-1)^2, & 0 \leq t \leq 1, 1 \leq x < 5, \\ 161 + 2500(x-5)^3, & 0 \leq t \leq 1, 5 \leq x < 6, \\ 2661 + 4(x-6)^2, & 0 \leq t \leq 1, x \geq 6. \end{cases}$$

Then (1) has at least two positive solutions.

Proof. Let $M = 3$, $d = 1$, $a = 5$, $b = 24$, $\delta = \frac{1}{4}$. Clearly, (H_1) – (H_3) hold, and $a, b, d > 0$ satisfy:

$$0 < \frac{3}{8} \Phi^{-1} \left[\Phi(8) + \frac{3}{8} \right] + 1 \leq 5 < 6 < 24.$$

After some simple calculation, we have

$$f(t, x) \geq 1 \text{ for } (t, x) \in \left[\frac{1}{8}, \frac{7}{8}\right] \times [1, 24],$$

$$f(t, x) < 200 \text{ for } (t, x) \in [0, 1] \times [0, 5],$$

$$f(t, y) \geq 2307 \text{ for } (t, x) \in [\frac{1}{4}, \frac{3}{4}] \times [6, 24],$$

$$f(t, x) \leq 12\,800 \text{ for } (t, x) \in [0, \frac{1}{8}] \times [0, 24].$$

Thus, by an application of Theorem 2, we get that problem (1) has at least two positive solutions. \square

References

- [1] R.P. Agarwal, D. O'Regan, P.J.Y. Wong, *Positive Solutions of Differential, Difference and Integral Equations*, Kluwer Academic publishers, Dordrecht, 1999.
- [2] R.I. Avery, Existence of multiple positive solutions to a conjugate boundary value problem, *MRS Hot-Line* 2 (1998) 1–6.
- [3] R.I. Avery, A generalization of the Leggett–Williams fixed point theorem, *Math. Sci. Res. Hot-Line* 3 (1999) 9–14.
- [4] R.I. Avery, Chuan Jen Chyan, J. Henderson, Twin solutions of boundary value problems for ordinary differential equations and finite difference equations, *Comput. Math. Appl.* 42(2001) 695–704.
- [5] R.I. Avery, J. Henderson, Three symmetric positive solutions for a second order boundary value problem, *Appl. Math. Lett.* 13 (2000) 1–7.
- [6] K. Deimling, *Nonlinear Functional Analysis*, Springer, New York, 1985.
- [7] J. Dugundji, An extension of Tietze theorem, *Pacific J. Math.* 1 (1951) 353–367.
- [8] L.H. Erbe, S. Hu, H. Wang, Multiple positive solutions of some boundary value problems, *J. Math. Anal. Appl.* 184 (1994) 640–648.
- [9] L.H. Erbe, H. Wang, On the existence of positive solutions of ordinary differential equations, *Proc. Amer. Math. Soc.* 120 (1994) 743–748.
- [10] D. Guo, V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, Academic Press, San Diego, 1988.
- [11] Guo Yanping, Ge Weigao, Gao Ying, Twin positive solutions for higher order m-point boundary value problems with sign changing nonlinearities, *Appl. Math. Comput.*, 146 (2003) 299–311.
- [12] J. Henderson, H.B. Thompson, Multiple symmetric positive solutions for a second-order boundary value problem, *Proc. Amer. Math. Soc.* 128 (2000) 2373–2379.
- [13] R.W. Leggett, L.R. Williams, Multiple positive fixed points of nonlinear operators on ordered Banach spaces, *Indiana Univ. Math. J.* 28 (1979) 673–688.
- [14] Wang Junyu, The existence of positive solutions for the one-dimensional p-Laplacian, *Proc. Amer. Math. Soc.* 125(1997) 2275–2283.